



020-222^{Q&As}

ACSM Health/Fitness Instructor Exam

Pass ACSM 020-222 Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/020-222.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by ACSM
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers





QUESTION 1

A client's health screening should be administered before

- A. Any contact with the client.
- B. Any physical activity by the client at your facility.
- C. Fitness assessment or programming.
- D. The initial "walk-through" showing of a facility.

Correct Answer: B

QUESTION 2

Category 1 medications include all of the following EXCEPT

- A. Lidocaine.
- B. Oxygen.
- C. Xylocaine.
- D. Epinephrine.

Correct Answer: C

QUESTION 3

A client must be given specific instructions for the days preceding a fitness assessment. Which of the following is NOT a necessary instruction to a client for a fitness assessment?

- A. Men and women should avoid liquids for 12 hours before the test.
- B. Clients should be instructed to avoid alcohol, tobacco products, or caffeine at least 3 hours before the test.
- C. Clients should avoid strenuous exercise or physical activity on the day of the test.
- D. Men and women should be instructed to get an adequate amount of sleep the night before the assessment.

Correct Answer: A

QUESTION 4

Which of the following would most appropriately assess a previously sedentary, 40-year-old female client's muscular strength?

- A. Using a 30-pound (18-kg) barbell to perform biceps curls to fatigue.



- B. Holding a handgrip dynamometer at 15 pounds (7 kg) to fatigue.
- C. Performing modified curl-ups to fatigue.
- D. Using a 5-pound (2.2-kg) dumbbell to perform multiple sets of biceps curls to fatigue.

Correct Answer: A

QUESTION 5

Complaints of pain in the chest with associated pain radiating down the left arm may be signs of

- A. Cardiac crisis.
- B. Hypotension.
- C. Seizure.
- D. Heartburn.

Correct Answer: A

[Latest 020-222 Dumps](#)

[020-222 Practice Test](#)

[020-222 Study Guide](#)