



AAPC-CPC^{Q&As}

Certified Professional Coder (CPC) Exam

Pass AAPC AAPC-CPC Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/aapc-cpc.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by AAPC
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





QUESTION 1

A patient suffers a ruptured infrarenal abdominal aortic aneurysm requiring emergent endovascular repair. An aorto-aortic tube endograft is positioned in the aorta and a balloon dilation is performed at the proximal and distal seal zones of the endograft. The balloon angioplasty is performed for endoleak treatment.

What CPT?code does the vascular surgeon use to report the procedure?

- A. 34702
- B. 34701
- C. 34707
- D. 34708

Correct Answer: A

QUESTION 2

A mother brings her 2-year-old son to the pediatrician's office because he stuck a bead up his left nostril. The pediatrician uses a nasal decongestant to open the blocked nostril and removes the bead with nasal forceps. What CPT?coding is reported?

- A. 30210-50
- B. 30210
- C. 30300
- D. 30300-50

Correct Answer: C

QUESTION 3

A patient with three thyroid nodules is seen for an FNA biopsy. Using ultrasonic guidance, the provider inserts a 25-gauge needle into each nodule. Nodular tissue is aspirated and sent to pathology. What CPT?coding reported?

- A. 10005, 10006 x 2, 76942
- B. 10006 x 3
- C. 10005, 10006 x 2
- D. 10021, 10004 x 2, 76942

Correct Answer: C



QUESTION 4

A 42-year-old with chronic left trochanteric bursitis is scheduled to receive an injection at the Pain Clinic. A 22-gauge spinal needle is introduced into the trochanteric bursa under ultrasonic guidance, and a total volume of 8 cc of normal saline and 40 mg of Kenalog was injected.

What CPT?code should be reported for the surgical procedure?

- A. 20610-LT
- B. 20611-LT, 76942
- C. 20611-LT
- D. 20610-LT, 76942

Correct Answer: C

QUESTION 5

View MR 002395 MR 002395 Operative Report Pre-operative Diagnosis: Acute rotator cuff tear Post-operative Diagnosis: Acute rotator cuff tear, synovitis Procedures: 1) Rotator cuff repair 2) Biceps Tenodesis 3) Claviculectomy 4) Coracoacromial ligament release Indication: Rotator cuff injury of a 32-year-old male, sustained while playing soccer. Findings: Complete tear of the right rotator cuff, synovitis, impingement. Procedure: The patient was prepared for surgery and placed in left lateral decubitus position. Standard posterior arthroscopy portals were made followed by an anterior- superior portal. Diagnostic arthroscopy was performed. Significant

synovitis was carefully debrided. There was a full-thickness upper 3rd subscapularis tear, which was repaired. The lesser tuberosity was debrided back to bleeding healthy bone and a Mitek 4.5 mm helix anchor was placed in the lesser tuberosity. Sutures were passed through the subcapulans in a combination of horizontal mattress and simple interrupted fashion and then tied. There was a partial-thickness tearing of the long head of the biceps. The biceps were released and then anchored in the intertubercular groove with a screw. There was a large anterior acromial spur with subacromial impingement. A CA ligament was released and acromioplasty was performed. Attention was then directed to the

supraspinatus tendon tear. The tear was V-shaped and measured approximately 2.5 cm from anterior to posterior. Two Smith and Nephew PEEK anchors were used for the medial row utilizing Healicoil anchors. Side-to-side stitches were placed. One set of suture tape from each of the medial anchors was then placed through a laterally placed Mitek helix PEEK knotless anchor which was fully inserted after tensioning the tapes. A solid repair was obtained. Next there were severe degenerative changes at the AC joint of approximately 8 to 10 mm. The distal clavicle was resected taking care to preserve the superior AC joint capsule. The shoulder was thoroughly lavaged. The instruments were removed and the incisions were closed in routine fashion. Sterile dressing was applied. The patient was transferred to recovery in stable condition.

What CPT?coding is reported for this case?

- A. 29827, 29828-51, 29824-51, 29826
- B. 29827, 29824-51, 29826-51
- C. 29827, 29828-51, 29824-51, 29826, 29805-59
- D. 29827, 29824-51, 29826-51, 29805-59

Correct Answer: A



VCE & PDF

PassApply.com

<https://www.passapply.com/aapc-cpc.html>

2024 Latest passapply AAPC-CPC PDF and VCE dumps Download

[AAPC-CPC Study Guide](#)

[AAPC-CPC Exam
Questions](#)

[AAPC-CPC Braindumps](#)