



ACLS^{Q&As}

Advanced Cardiac Life Support

Pass Test Prep ACLS Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/acls.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





QUESTION 1


FILL BLANK

What is the most important goal of ACLS?

A. Restoration of coronary perfusion with subsequent spontaneous cardiac rhythm and preservation of brain function.

Correct Answer: A

QUESTION 2



Identify the rhythm by selecting the best single answer:


- A. Normal sinus rhythm
- B. Sinus tachycardia
- C. Sinus bradycardia
- D. Reentry supraventricular tachycardia
- E. First-degree AV Block
- F. Second-degree AV Block (Mobitz 1 Wenckebach)
- G. Second-degree AV Block (Mobitz II Block)
- H. Third-degree AV Block
- I. Atrial fibrillation
- J. Atrial flutter
- K. Monomorphic ventricular tachycardia
- L. Polymorphic ventricular tachycardia



- M. Coarse ventricular fibrillation
- N. Fine ventricular fibrillation
- O. Agonal rhythm/asystole
- P. Pulseless electrical activity

Correct Answer: C

QUESTION 3



A patient has been resuscitated from cardiac arrest and is being prepared for transport. She is incubated and is receiving 100% oxygen. During the resuscitation she received 2 doses of epinephrine 1 mg, atropine 1 mg, and lidocaine 100 mg IV. You now observe the above rhythm on the cardiac monitor. The rhythm abnormality is becoming more frequent and increasing in number. You should order:

- A. Give amiodarone 300 mg IV, start infusion
- B. Give lidocaine 1 to 1.5 mg IV
- C. Repeat epinephrine 1 mg IV
- D. Give lidocaine 0.5 to 0.75 mg/kg IV, start lidocaine infusion
- E. Give amiodarone 150 mg IV, start infusion

Correct Answer: D

QUESTION 4




This patient suddenly collapsed and is poorly responsive. The patient has a weak carotid pulse. A cardiac monitor, oxygen, and an intravenous line have been initiated. The code cart with all drugs and pF transcutaneous pacer is immediately available. Next you would

- A. Initiate dopamine at 10 to 20 ug/kg per minute and titrate heart rate
- B. Give atropine 1 mg IV up to a total dose of 3 mg
- C. Initiate epinephrine at 2 to 10 ug per minute and titrate heart rate
- D. Initiate dopamine at 2 to 10 ug/kg per minute and titrate heart rate
- E. Begin transcutaneous pacing

Correct Answer: E

QUESTION 5

A patient with a myocardial infarction develops the following rhythm and loses consciousness. The patient is pulseless and not breathing. What is the preferred immediate treatment?



- A. administer bretylium
- B. administer lidocaine
- C. call for help and deliver a synchronized shock
- D. call for help and deliver an unsynchronized shock

Correct Answer: D



VCE & PDF

PassApply.com

<https://www.passapply.com/acls.html>

2024 Latest passapply ACLS PDF and VCE dumps Download

[ACLS Study Guide](#)

[ACLS Exam Questions](#)

[ACLS Braindumps](#)