



HAAD-RN^{Q&As}

HAAD Licensure Examination for Registered Nurses

Pass HAAD HAAD-RN Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/haad-rn.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by HAAD
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





QUESTION 1

A newborn infant is assessed using the Apgar assessment tool and scores 6. The infant has a heart rate of 95, slow and irregular respiratory effort, and some flexion of extremities. The infant is pink, but has a weak cry. The nurse should know that this Apgar score along with the additional symptoms indicates the neonate is:

- A. Functioning normally
- B. Needing immediate life-sustaining measures
- C. Needing special assistance
- D. Needing to be warmed

Correct Answer: C

QUESTION 2

A patient with a history of angina pectoris arrives in emergency complaining of headache, visual disturbances and feeling dizzy. Your nursing assessment also notes he looks flushed, is perspiring profusely and is experiencing palpitations. You should suspect:

- A. An overdose of sublingual nitroglycerin
- B. The onset of a myocardial infarction
- C. The patient has been over exercising
- D. The beginning of a severe chest infection

Correct Answer: A

QUESTION 3

A newborn is diagnosed with ventricular septal defect. The baby is discharged with a prescription for digoxin syrup 20 micrograms bid. The bottle of digoxin is labeled as 0.05 mg/ml. The nurse should teach the mother to administer on each dose:

- A. 0.1 ml
- B. 0.2 ml
- C. 0.4 ml
- D. 0.8 ml

Correct Answer: C

QUESTION 4



A 3-month-old infant is admitted with a diagnosis of ventricular septal defect. The physical assessment for this infant would reveal:

- A. High pitched cry
- B. Harsh heart murmur
- C. Bradycardia
- D. Hypertension

Correct Answer: B

QUESTION 5

Symptoms of alcohol withdrawal include:

- A. Euphoria, hyperactivity and insomnia
- B. Depression, suicidal ideation and hypersomnia
- C. Diaphoresis, nausea and vomiting and tremors
- D. Unsteady gait, nystagmus and profound disorientation

Correct Answer: C

[HAAD-RN PDF Dumps](#)

[HAAD-RN Practice Test](#)

[HAAD-RN Brainsdumps](#)