



# MCQS<sup>Q&As</sup>

Multiple-choice questions for general practitioner (GP) Doctor

## Pass Test Prep MCQS Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/mcqs.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep  
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





### QUESTION 1

Chronic pancreatitis, investigation of choice?

- A. ERCP
- B. MRCP
- C. MRI
- D. CECT

Correct Answer: A

---

### QUESTION 2

A 26-year-old male has been operated for abdominal trauma and splenectomy was done. On the 3rd post-op day the developed acute abdominal pain and distention in the upper abdominal area with hypotension. On insertion of ryle's tubes, 2L of coffee ground fluid was aspirated.

What is the most probable diagnosis?

- A. Reactionary hemorrhage
- B. DVT
- C. Acute gastric dilatation
- D. Left lower lobe atelectasis
- E. Subphrenic abscess

Correct Answer: D

Patients with blood pressures (BP) > 180/120 mm Hg need immediate aggressive treatment if they have impending or progressive end-organ damage. Those without end-organ damage need to adjust or reinstate their regimens, but do not need immediate aggressive treatment.

---

### QUESTION 3

Pregnant woman with UTI which is the best antibiotics to be given if she has no allergy?

- A. Tetracycline
- B. Aminoglycoside
- C. Sulfa Trimethoprim
- D. Ampicillin
- E. Nitrofurantoin



Correct Answer: E

Urinary tract infections are common during pregnancy, and the most common causative organism is Escherichia coli. Asymptomatic bacteriuria can lead to the development of cystitis or pyelonephritis. All pregnant women should be screened for bacteriuria and subsequently treated with antibiotics such as nitrofurantoin, sulfisoxazole or cephalexin. Ampicillin should no longer be used in the treatment of asymptomatic bacteriuria because of high rates of resistance. Pyelonephritis can be a life-threatening illness, with increased risk of perinatal and neonatal morbidity. Recurrent infections are common during pregnancy and require prophylactic treatment. Pregnant women with urinary group B streptococcal infection should be treated and should receive intrapartum prophylactic therapy.

---

#### QUESTION 4

Patient with major depression what is the first line treatment?

A. Tricyclic antidepressant

B. SSRIs

C. MAOI

Correct Answer: B

A SSRIs are the initial antidepressants of choice for uncomplicated depression because of their minimal anti-cholinergic effects

---

#### QUESTION 5

The tumor, which may occur in the residual breast or overlying skin following wide local excision and radiotherapy for mammary carcinoma, is:

A. Squamous cell carcinoma

B. Leiomyosarcoma

C. Angiosarcoma

D. Basal cell carcinoma

Correct Answer: C

[Latest MCQS Dumps](#)

[MCQS PDF Dumps](#)

[MCQS VCE Dumps](#)