



# NAPLEX<sup>Q&As</sup>

North American Pharmacist Licensure Examination

## Pass Test Prep NAPLEX Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/naplex.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep  
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers





### QUESTION 1

A patient takes 1gm of Calcium Carbonate salt three times a day. How much elemental calcium, in grams, is he getting in 24hrs? (MW of Ca: 40.078 g/mol, MW of CaCO<sub>3</sub>: 100.087 g/mol)

- A. 3 g
- B. 1.8g
- C. 1.2g
- D. 0.8gm
- E. 1.8mg

Correct Answer: C

---

### QUESTION 2

Mesna is typically administered alongside which of these chemotherapeutic agents?

- A. Busulfan
- B. Paclitaxel
- C. Ifosfamide
- D. Doxorubicin

Correct Answer: C

Mesna is administered to patients taking either ifosfamide or cyclophosphamide to conjugate toxic acrolein and therefore prevent hemorrhagic cystitis (bladder bleeding).

---

### QUESTION 3

Which of the following antidiabetic medication works by inhibiting carbohydrate breakdown?

- A. Acarbose
- B. Metformin
- C. Dapagliflozin
- D. Pioglitazone
- E. Sitagliptin

Correct Answer: A



Acarbose is an alpha glucosidase inhibitor that inhibits carbohydrate breakdown. Metformin is a biguanide that decreases hepatic glucose production. Dapagliflozin is a SGLT2 inhibitor to decrease glucose reabsorption in the kidney. Pioglitazone is a TZD that increases insulin sensitivity. Sitagliptin is a DPP-4 inhibitor that works on incretins/increase insulin secretion/decrease glucagon secretion.

---

**QUESTION 4**

What is the active ingredient found in the medicine Adalat?

- A. Nifedipine
- B. Adalimumab
- C. Digoxin
- D. Simvastatin

Correct Answer: A

---

**QUESTION 5**

A fasting blood glucose level greater than what value is an indicator of type 2 diabetes?

- A. 5 mmol/L
- B. 6 mmol/L
- C. 7 mmol/L
- D. 8 mmol/L
- E. 9 mmol/L

Correct Answer: C

A fasting blood glucose level of greater than 7 mmol/L (or greater than 126mg/dL) is an indicator of type 2 diabetes.

[Latest NAPLEX Dumps](#)

[NAPLEX VCE Dumps](#)

[NAPLEX Exam Questions](#)