



# RPFT<sup>Q&As</sup>

Registry Examination for Advanced Pulmonary Function Technologists

## Pass Test Prep RPFT Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/rpft.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep  
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





### QUESTION 1

An isothermal lung analog has the following specifications:

5-L glass bottle 2000 g of steel wool Density of steel wool 5 g/cm<sup>3</sup> Volume of connectors and bulb 100 ml Repeated FRC measurements in the plethysmograph result in a volume of 5.5 L.

Which of the following should a pulmonary function technologist do?

- A. Remove 500 g of steel wool from the lung analog
- B. Take the plethysmograph out of service
- C. Recalculate the BTPS correction
- D. Begin a biologic control program

Correct Answer: D

---

### QUESTION 2

During an exercise study, the RER will equal the RQ only when the patient is at

- A. Steady state
- B. Peak exercise
- C. The anaerobic threshold
- D. Rest

Correct Answer: A

---

### QUESTION 3

A pulmonary function laboratory routinely tests five subjects as biologic controls. Acceptable instrument performance over time is indicated when measurements

- A. For each individual are within 2 standard deviations of the mean of previous measurements for each individual.
- B. For an individual are within 2 standard deviations of the mean of those for the group.
- C. Of the group as a whole agree with those predicted by linear regression for changes in age.
- D. Are within the 95% confidence interval predicted for an individual being tested.

Correct Answer: D

---

### QUESTION 4



Which of the following is a suitable policy for following Standard Precautions in a pulmonary function laboratory?

- A. Eye protection is required when obtaining ABGs from patients with hepatitis.
- B. Reusable mouthpieces should be disposed when a patient has a history of tuberculosis.
- C. Gloves are optional when obtaining arterial blood samples using a kit
- D. Reusable mouthpieces should be disinfected between each patient.

Correct Answer: B

---

#### QUESTION 5

A patient will undergo bronchial provocation testing with methacholine chloride (Provocholine). Baseline spirometry reveals the FVC is 4.00 L and FEV1 is 3.00 L. The provocative dose 20% (PD20) can be calculated when which of the following measurements occur?

- A. The FVC falls below 3.20 L, or the FEV1 falls below 2.40 L.
- B. The FEV<sub>1</sub> falls below 2.40 L, regardless of change in FVC.
- C. The FVC falls below 3.20 L and the FEV1 falls below 2.40 L.
- D. The ratio FEV<sub>1</sub>/FVC falls below 0.8.

Correct Answer: B

[Latest RPFT Dumps](#)

[RPFT VCE Dumps](#)

[RPFT Exam Questions](#)