



# HAAD-RN<sup>Q&As</sup>

HAAD Licensure Examination for Registered Nurses

**Pass HAAD HAAD-RN Exam with 100% Guarantee**

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.passapply.com/haad-rn.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by HAAD  
Official Exam Center

-  **Instant Download** After Purchase
-  **100% Money Back** Guarantee
-  **365 Days** Free Update
-  **800,000+** Satisfied Customers





#### QUESTION 1

Collection urine bag should be emptied as necessary and at least every 8 - 9 hours to prevent:

- A. Pooling of urine in the tube
- B. Reflux of urine into the bladder
- C. Pulling on catheter
- D. Bacterial contamination

Correct Answer: D

---

#### QUESTION 2

The urinary catheter is kept securely in the bladder by:

- A. Taping the urinary catheter to the leg
- B. Securing catheter and collection bag connections
- C. Inflating the balloon of the catheter
- D. Anchoring the catheter bag to the bed

Correct Answer: C

---

#### QUESTION 3

Which of the following indicates the nurse is engaging in a therapeutic nurse-patient relationship?

- A. The nurse establishes a relationship that is mutually beneficial
- B. The nurse demonstrates sympathetic feelings toward the patient
- C. The nurse commits to helping the patient find ways to help self
- D. The nurse utilizes therapeutic touch to convey acceptance of the patient

Correct Answer: C

---

#### QUESTION 4

A nursing measure that is helpful in communicating with a hearing impaired patient is to:

- A. Use simple sentences
- B. Talk while close to patient's ear



- C. Raise the voice
- D. Write out all questions and responses

Correct Answer: A

---

#### QUESTION 5

Which of the following interventions should the nurse implement if a patient complains of cramps while irrigating the colostomy?

- A. Reduce the flow of solution
- B. Have the patient sit up in bed
- C. Remove the irrigation tube
- D. Insert the tube further into the colon

Correct Answer: A

---

#### QUESTION 6

To ensure safe practice during defibrillation, the nurse must:

- A. Cover paddles with electrode gel
- B. Avoid touching the patient's bed
- C. Remove paddles after the shock
- D. Synchronize prior to shock delivery

Correct Answer: B

---

#### QUESTION 7

While assessing a patient, the nurse learns that he has a history of allergic rhinitis, asthma, and multiple food allergies. The nurse must:

- A. Be alert to hypersensitivity response to the prescribed medications
- B. Encourages the patient to carry an epinephrine kit in case of an allergic reaction
- C. Advise the patient to use aspirin in case of febrile illnesses
- D. Admit the patient to a single room with limited exposure to health care personnel

Correct Answer: A

---



### QUESTION 8

The patient is receiving piperacillin (Pipril) I.V. Which of the following should the nurse consider when administering the medication to the patient?

- A. Assess for Stevens-Johnson Syndrome
- B. Be alert for the possibility of hypersensitivity
- C. Watch for signs of increasing severity of infection
- D. Instruct patient to increase the intake of protein-rich foods

Correct Answer: B

---

### QUESTION 9

A patient with deep partial-thickness and full-thickness burns of the face and chest is admitted to the emergency department. The nurse must be particularly alert for:

- A. Paralytic ileus
- B. Respiratory distress
- C. Severity of pain
- D. Strong burn odor

Correct Answer: B

---

### QUESTION 10

When caring for a patient with hepatic encephalopathy the nurse may carry out the following orders: give enemas, provide a low protein diet, and limit physical activities.

These measures are performed to:

- A. Minimize edema
- B. Decrease portal pressure
- C. Reduce hyperkalemia
- D. Decrease serum ammonia

Correct Answer: D

---

### QUESTION 11

The patient has a nursing diagnosis of altered cerebral tissue perfusion related to cerebral edema. An appropriate



nursing intervention for this problem is to:

- A. Elevate the head of the bed 30 degrees
- B. Provide a position of comfort with knee flexion
- C. Provide uninterrupted periods of rest
- D. Ensure adequate hydration with mannitol

Correct Answer: A

---

#### QUESTION 12

The rationale for having the patient void before an abdominal paracentesis procedure is to:

- A. Minimize discomfort
- B. Avoid abdominal distention
- C. Prevent bladder puncture
- D. Reduce infection rate

Correct Answer: C

---

#### QUESTION 13

When the nurse is caring for a patient placed on droplet precautions, the nurse should:

- A. Have the patient wear a high-efficiency particulate air (HEPA) mask
- B. Wear a surgical mask when standing within 3 feet (1 meter) of the patient
- C. Assign the patient to a room with monitored negative air pressure
- D. Apply a disposable gown when entering the patient's room

Correct Answer: B

---

#### QUESTION 14

To remove soft contact lenses from the eyes of an unconscious patient the nurse should:

- A. Uses a small suction cup placed on the lenses
- B. Pinches the lens off the eye then slides it off the cornea
- C. Lifts the lenses with a dry cotton ball that adheres to the lenses
- D. Tenses the lateral canthus while stimulating a blink reflex by the patient



Correct Answer: B

---

#### QUESTION 15

A construction worker was brought to the emergency department and admitted with the diagnosis of heat stroke due to strenuous physical activity during hot weather conditions.

Which action should the nurse take?

- A. Immediately immerse the patient in cold water to reduce the patient's temperature
- B. Administer an antipyretic such as aspirin or acetaminophen
- C. Place ice packs to the neck, axillae, scalp and groin
- D. Encourage foods and oral fluids that contain carbohydrates and electrolytes

Correct Answer: C

[HAAD-RN PDF Dumps](#)

[HAAD-RN Exam Questions](#)

[HAAD-RN Braindumps](#)